

IPSC-Based PDAC Models and Immunotherapies

Subjects: [Oncology](#) | [Cell Biology](#)

Contributor: Ricki T. Krog

Advances in the treatment of pancreatic ductal adenocarcinoma (PDAC) using neoadjuvant chemoradiotherapy, chemotherapy, and immunotherapy have had minimal impact on the overall survival of patients. A general lack of immunogenic features and a complex tumor microenvironment (TME) are likely culprits for therapy refractoriness in PDAC. Induced pluripotent stem cells (iPSCs) should be explored as a means to advance the treatment options for PDAC, by providing representative in vitro models of pancreatic cancer development. In addition, iPSCs could be used for tailor-made cellular immunotherapies or as a source of tumor-associated antigens in the context of vaccination.

pancreatic ductal adenocarcinoma

PDAC

induced pluripotent stem cells

iPSCs

cancer models

cancer vaccine

cancer therapy

immunotherapy

microenvironment

1. Introduction

Pancreatic ductal adenocarcinoma (PDAC) is the most common type of pancreatic cancer, accounting for more than 90% of all cases ^[1]. PDAC has an extremely poor survival rate, of approximately 9% at five years, which has remained stable in the last decades despite improvements in the treatment of other cancer types ^{[1][2]}. Alarming, pancreatic cancer is predicted to become the second most common cause of cancers by 2030 in the United States ^[3]. For a long time, the standard treatment of PDAC consisted of chemotherapy or radiotherapy as a monotherapy. In recent years, this has changed towards slightly more efficient combined neoadjuvant treatments followed by surgical resection. Surgery is the only curative option of treatment for PDAC patients, but it applies to fewer than 20% of patients that are diagnosed in the early stage of disease prior to locally advanced, borderline unresectable, or metastatic disease ^[4]. Limited knowledge of both clinical symptoms and reliable biomarkers for precancerous lesions, such as pancreatic intraepithelial neoplasm (PanIN) ^[5], intraductal papillary mucinous neoplasm (IPMN) (reviewed in ^[6]), and acinar-to-ductal metaplasia (ADM) ^{[7][8]}, are major obstacles for early diagnosis ^[9].

2. The Potential of Induced Pluripotent Stem Cells (iPSCs)

iPSCs provide excellent possibilities for the modeling of PDAC and serve as a unique source of tumor-associated antigens for whole-cell cancer vaccines and immune cells for adoptive cellular immunotherapies. Pluripotent stem cells hold stemness, which is defined as the ability to proliferate indefinitely while maintaining pluripotency. Embryonic stem cells (ESCs) are a classic example of this capability and are capable of differentiating into all of

the three embryonic germ layers, thus giving rise to all adult cell types [10][11]. Since the isolation of ESCs, scientists have mainly focused on their extraordinary potential for (personalized) regenerative medicine. Takahashi and Yamanaka demonstrated in 2006 that defined culture conditions, including the exogenous supply of four transcription factors (Yamanaka factors) MYC, OCT3/4, SOX2, and KLF4, were capable of reprogramming differentiated mouse embryonic and adult fibroblast cells into cells with characteristics of ESCs [12]. These dedifferentiated pluripotent cells were termed induced pluripotent stem cells (iPSCs). This discovery was immediately followed by the induction of human iPSCs from terminally differentiated cells, and in subsequent years various methods have been developed to generate iPSCs from a variety of somatic cells (reviewed in [13][14] [15][16] [17][18][19][20][21][22][23][24][25][26]). These accomplishments led to the exploration of the potential of iPSCs for disease modeling and treatment of various diseases [15][16]. iPSCs share a number of characteristics with ESCs with almost identical gene expression and epigenetic status and possess the unique feature of stemness [27][28][29][30]. Furthermore, ethical issues associated with human ESCs are avoided; thus, iPSCs expand the range of applications in which stem cells can be exploited. These applications include basic cell biology, disease models, and drug discovery and screening. Furthermore, iPSCs provide the potential for new clinical applications and can serve as the basis for (off-the-shelf) cancer immunotherapies. The pros and cons of iPSCs related to cancer immunotherapies and models are listed in **Table 1**.

Table 1. Pros and cons of induced pluripotent stem cells (iPSCs) related to cancer immunotherapies and models.

	Pros +	Cons -
Models [31] [32] [33] [34] [35] [36] [37]	+ Insight into disease onset and development	- Risk of transcriptional and genetic alterations during iPSC induction
	+ Reliable cell source for multicellular models	- Heterogeneity within iPSC populations
Immunotherapeutic approaches [38] [39] [40] [41] [42] [43] [44] [45] [46] (ClinicalTrials.gov, Identifier: NCT03841110 jrct.niph.go.jp, Trial ID: jRCT2033200116)	+ iPSCs are immunogenic and express tumor-associated antigens	- Oncogenic potential of iPSCs reprogramming factors
	+ Reliable cell source for off-the-shelf cellular therapies	- Risk of transcriptional and genetic alterations during iPSC induction
	+ Large-scale cell production	- Heterogeneity within iPSC populations

3. iPSC-Based PDAC Models and Their Potential for Disease Modeling

Around 20% of PDAC patients in the United States are diagnosed with localized disease and are therefore eligible for surgical resection [2][4]. However, the majority of patients are diagnosed with locally advanced or metastasized disease at diagnosis, which leaves these patients with a poor survival rate [2]. Limited knowledge of both clinical symptoms and biomarkers in the early stages of PDAC are major obstacles for early disease stage diagnosis [9]. A number of genomic alterations have been associated with PDAC, but our understanding of their precise role in the onset and progression of disease is limited as the genomic studies associated with disease progression are sparse due to the lack of suitable models. For example, PDAC-derived xenografts have been established in immunocompromised mice by using tumor tissues or cell lines [47][48][49][50][51]. These models solely reflect the invasive stages of PDAC and are not suitable for studies on the onset and early stages of PDAC. Novel models providing a better understanding of the biological processes at the basis of tumorigenesis are essential to improve diagnostics. To address this, iPSCs can provide a source of cells that better reflect the early stages of malignant transformation in PDAC.

iPSC-derived cancer-initiating cells have previously been reported for the establishment of xenograft models that reflect the malignant transformation in PDAC [32][52]. Mouse iPSCs from healthy cells have been differentiated in a controlled manner into PDAC progenitor cells [32]. Xenograft models originating from these cells were able to give rise to precancerous lesions, including ADM and PanIN, as well as invasive PDAC. Exploiting a different approach, Kim et al. (2013) hypothesized that a subset of iPSCs induced from human PDAC cells would result in malignant iPSC lines, capable of undergoing early developmental stages of PDAC after engraftment into mice [34]. One of the generated iPSC cell lines carried a *KRAS*^{G12D} mutation and a deletion of *CDKN2A*. The oncogenic *KRAS* mutations are the most frequently detected oncogenic alteration in PDAC, being observed in >90% of patients [53][54][55]. *CDKN2A* is a tumor suppressor in PDAC and has been described as being inactivated in approximately 50% of patients [56][57]. Xenografts originating from the *KRAS*^{G12D} *CDKN2A*^{-/-} iPSC cell line gave rise to PanIN-like lesions followed by progression to invasive PDAC [34]. iPSC-based xenograft PDAC models originating from malignant cells demonstrate the potential of iPSCs to provide insights into PDAC onset and progression, including the identification of potential biomarkers for early diagnosis of PDAC. Another application where iPSCs might improve PDAC-modeling is the generation of iPSC-derived organoids containing different cell populations. iPSCs can be committed to a differentiation into the pancreatic exocrine lineage for the generation of acinar and ductal cells and, thus, provide great organoid-modeling possibilities for PDAC [31][33][58][59]. PDAC can develop from both acini and ducts, however knowledge on how these two cells of origin impact cell progression is scarce [60]. Two studies recently assessed how the PDAC oncogenes *KRAS* and *GNAS* individually affect the growth and progression of PDAC in vitro and in vivo after engraftment of iPSC-derived acinar and ductal organoids in immunocompromised mice [31][33]. Both *KRAS*^{G12D}-mutated acinar and ductal organoids displayed proliferation in vivo, although the more invasive lesions were generated from acinar organoids. Phenotypically, both oncogenic alterations caused IPMN-like lesions in vivo. Furthermore, PanIN lesions and different stages of PDAC-like tumor formation were observed in xenografts from *KRAS*^{G12D}-mutated ductal and acinar organoids. In vitro, *KRAS*^{G12D}-mutated ductal organoids displayed epithelial-to-mesenchymal transition (EMT), which have been suggested to

play a role in early tumor formation, metastasis, and chemoresistance in PDAC [61][62][63]. In contrast, GNAS^{R201C/H} induced cystic growth in vitro in ductal organoids and to a lesser extent in acinar organoids. These iPSC-derived models provide vital knowledge of the malignant potential of different oncogenes in PDAC. Furthermore, the models provide great opportunities for in-depth assessment of early-stage disease development and progression.

In addition to the above-mentioned applications of iPSCs for disease modeling, iPSCs can also be differentiated into non-malignant cells of the TME. This opens up avenues for the development of complex multicellular models to test therapeutic interventions. For example, TAMs are thought to play an important role in PDAC tumorigenesis and may constitute promising clinical targets [64]. Macrophage models for drug discovery have so far been dependent on a limited source of monocytes derived from PBMCs or animal bone marrow, which has limited the generation of models representative of tissue-resident macrophages (reviewed in [65]). Gutbier and colleagues established a method for controlled large-scale iPSC-derived tissue-resident-resembling macrophages for efficient drug screening and discovery [66]. Genetic manipulation of these iPSC-derived macrophages can be conducted to obtain the desired macrophage subtype. Additionally, cancer-initiating cells originating from iPSCs from healthy cells can also be differentiated into CAFs and vascular endothelial-like cells in vivo [35][36][37]. Particularly, CAFs have been implicated as important players in the tumorigenesis of PDAC (reviewed in [67]) [68][69]. CAFs constitute a promising therapeutic target in PDAC and several therapeutic strategies have been investigated preclinically and clinically (reviewed in [70][71]). The versatility of iPSCs to generate a variety of cells from the TME can support the development of models that include various cell types [72]. Additionally, the directed differentiation towards a cell line of interest shows the potential of iPSC-derived models for drug screening at the molecular level.

iPSC-based xenografts and organoids provide excellent innovative possibilities for the modeling of PDAC, especially to study precancerous lesions and the development of this disease. Furthermore, the potential of iPSCs as a source for a variety of cells provides an opportunity for the establishment of multicellular models that better represent the PDAC TME. However, iPSC-based PDAC models are still in the early phase and further research is needed to fully exploit their potential.

4. iPSCs as a Cell-Based Immunotherapy

PDAC is characterized by a low mutational burden and, consequently, a low amount of neoantigens are generated for spontaneous antitumor immune responses by the adaptive immune system [73]. Additionally, the highly immunosuppressive TME of PDAC further contributes to its poor immunogenic character. A classical way of stimulating a specific immune response is by antigen vaccination. Therapeutic cancer vaccines aim to stimulate antitumor immunity, e.g., by supporting the activation of cancer-specific CD8⁺ and CD4⁺ T cells. ESCs have been hypothesized to serve as an efficient source of antigens for cancer vaccines due to their immunogenicity and shared antigenic profile with cancer cells [74][75][76]. Similar to ESCs, iPSCs have an immunogenic potential and share antigens with PDAC cells, making them an attractive source of antigens for cancer vaccination (reviewed in [77]) [41][44][78]. Kooreman et al. have found that an iPSC-based cancer vaccine was capable of eliciting an immune response towards shared iPSC and cancer cell antigens in murine cancer models [41]. The vaccine consisted of autologous iPSCs to minimize alloimmunity and the toll-like receptor 9 (TLR9) agonist CpG, to enhance the

immunostimulatory properties of the vaccine. In murine models of breast, lung, and skin cancer, this iPSC-based cancer vaccine elicited a potent humoral and cell-mediated immune response sufficient to prevent or limit tumor growth *in vivo* without any observed associated adverse effects [41]. In a mouse model of PDAC, the same vaccine induced protective immunity characterized by the expansion of effector and memory CD8⁺ T cells, CD4⁺ T cells (excluding T_{reg} cells), and B cells, while reducing the amount of T_{reg} cells in tumor-draining lymph nodes [44]. Furthermore, several cancer-signature peptide antigens, containing previously experimentally reported T cell epitopes, were identified in the vaccine *in silico*; this suggests the possibility for expansion of PDAC antigen-specific effector T cells [44]. This iPSC-based vaccine was proposed as a promising tool to be employed in an adjuvant context, in combination with conventional therapies [41]. Furthermore, this type of vaccination could hold promise in a prophylactic or (neo)adjuvant setting to stimulate the immune system in combination with other immunotherapies, such as immune checkpoint inhibitors, adoptive transfer of primed autologous T cells, chimeric antigen receptor (CAR) cells, or modulators of the immunosuppressive TME, by targeting the WNT-signaling pathway (reviewed in [79] [80]).

Prophylactic vaccines can play a role in the prevention of cancers caused by viruses, e.g., human papillomaviruses (HPVs) and the hepatitis B virus (HBV). To date, no prophylactic vaccines aiming at preventing non-viral-related cancers have been approved. Lu et al. (2020) reported an iPSC-based autologous prophylactic cancer vaccination regimen that was evaluated using the KPC mouse model of PDAC with spontaneous tumor development [43]. The PDAC driver mutations Kras^{G12D} and p53^{R172H} were introduced in murine iPSCs derived from healthy cells followed by controlled differentiation into PDAC progenitor cells. These cells were antigenically comparable to PDAC cells from the KPC mice and, therefore, serve as a good repertoire of PDAC-expressing antigens. The iPSC-induced PDAC progenitor cells were infected with a non-replicating oncolytic virus to enhance the immunogenicity of the final vaccine formulation. Importantly, the immunogenicity of the vaccine demonstrated a clinical impact by delaying tumor development and prolonging survival of the KPC mice vaccinated before tumor development. However, the vaccine failed to provide complete protection from tumor development in the mice. Upon vaccination, CD8⁺ T-cell infiltration increased in the PDAC TME and an accumulation of central memory T cells was observed in the secondary lymph nodes. Furthermore, splenocytes from vaccinated mice, prior to tumor development, showed enhanced production of the proinflammatory cytokine IFN γ after *ex-vivo* challenge with tumor-cell lines derived from the corresponding model. This iPSC-based autologous prophylactic cancer vaccination regimen demonstrates promising results in a mouse model of PDAC and further studies will clarify whether iPSC-based prophylactic PDAC vaccines have a potential for the prevention of pancreatic malignancies in at-risk individuals.

The first clinical use of iPSC-based cancer vaccines is yet to be seen, but the above-mentioned preclinical studies demonstrate the potential of these vaccines to elicit anti-PDAC immune responses. In addition to PDAC therapy, iPSC-based antitumor vaccination could potentially serve as a promising universal approach in a broad spectrum of cancer types, including mesothelioma, breast cancer, and melanoma (reviewed in [81] [41][82][83][84]). Tumorigenic properties of iPSCs necessitate lethal irradiation of the iPSCs prior to injection into patients to avoid a potential risk of tumor formation (reviewed in [85][86] [41][44][87][88]). Additionally, care must be taken to avoid activity of remnant transcription factors used for the induction of the iPSCs (**Figure 1**). Three of the four Yamanaka factors, MYC,

OCT3/4, and SOX2, are proto-oncogenes and are involved in tumorigenesis [76][89][90][91][92][93]; these transcription factors have been extensively reviewed elsewhere [94][95][96]. Efficient screening and purification must be carried out to secure a safe vaccine formulation before clinical implementation.




Figure 1. Risks associated with the induction of induced pluripotent stem cells (iPSCs) and the generation of a homogenous non-tumorigenic population of iPSC-derived immune cells. **(A)** Transcriptional alterations (aberrant DNA methylation) can occur in some iPSC clones during the induction of iPSCs leading to clones with a lower differential potential. This heterogenous iPSC induction, due to an improper epigenetic status of some iPSC clones, constitutes a major limitation in the induction of a homogenous population of iPSCs, which should be addressed by optimized reprogramming protocols; **(B)** Induction of iPSCs includes the risk of genetic alterations by the

introduction of single nucleotide variants (SNVs) and/or chromosomal aberrations. The mutation rate has been demonstrated to be around 10 times lower for iPSCs compared to somatic cells. However, genetic alterations will occur and presents an issue in the induction of a homogenous population of iPSCs for further applications. The risk of genetic alterations, influencing the phenotype of a subpopulation of iPSCs, highlights the importance of efficient screening and purification of the induced iPSCs; **(C)** Induction of iPSCs is carried out with the use of reprogramming factors with a potential tumorigenic potential. This necessitates the efficient removal of the reprogramming factors from the final iPSC population prior to clinical applications; **(D)** Remnant undifferentiated iPSCs constitute a potential risk for tumor formation in patients. Efficient purification and screening must be conducted to avoid undifferentiated iPSCs in the final population of iPSC-derived immune cells for clinical applications.

5. iPSC-Derived Immune Cells for Cancer Immunotherapies

Current approaches focusing on T cell immunotherapies, such as autologous T cell transfer, engineered T cell receptor (TCR) T cells, and CAR T cells, typically require autologous cell manufacturing for each individual patient. Therefore, there is an unmet need for innovative cell sources to broaden the application of cellular immunotherapies [97]. iPSCs can be a permanent source of various immune cells, which can be genetically modified for optimal therapeutic features. Despite being in the early stages, iPSC-derived NK cells (ClinicalTrials.gov, Identifier: NCT03841110) and NKT cells (jrct.niph.go.jp, Trial ID: jRCT2033200116) are currently being clinically investigated in patients with advanced solid tumors. It is worth noting that PDAC patients will be included for monotherapy with the iPSC-derived NK cells or as a combinatorial therapy with immune checkpoint inhibitors. With the increasing availability of promising preclinical data, it is plausible that additional clinical trials will be initiated in the near future. However, before clinical implementation, all safety issues must be thoroughly addressed (discussed in the original paper). Taken together, iPSCs serve as a promising approach for a novel source of a variety of immune cells for adoptive cellular immunotherapies.

References

1. Wild, C.P.; Weiderpass, E.; Stewart, B.W. Cancer Research for Cancer Prevention World Cancer Report; World Health Organization: Lyon, France, 2020.
2. National Cancer Institute. SEER Cancer Stat Facts: Pancreatic Cancer. Available online: <https://seer.cancer.gov/statfacts/html/pancreas.html> (accessed on 1 July 2021).
3. Rahib, L.; Smith, B.D.; Aizenberg, R.; Rosenzweig, A.B.; Fleshman, J.M.; Matrisian, L.M. Projecting Cancer Incidence and Deaths to 2030: The Unexpected Burden of Thyroid, Liver, and Pancreas Cancers in the United States. *Cancer Res.* 2014, 74, 2913–2921.
4. Siegel, R.L.; Miller, K.D.; Fuchs, H.E.; Jemal, A. Cancer Statistics, 2021. *CA A Cancer J. Clin.* 2021, 71, 7–33.

5. Hruban, R.H.; Adsay, N.V.; Albores–Saavedra, J.; Compton, C.; Garrett, E.S.; Goodman, S.N.; Kern, S.E.; Klimstra, D.S.; Klöppel, G.; Longnecker, D.S.; et al. Pancreatic Intraepithelial Neoplasia. *Am. J. Surg. Pathol.* 2001, 25, 579–586.
6. Patra, K.C.; Bardeesy, N.; Mizukami, Y. Diversity of Precursor Lesions For Pancreatic Cancer: The Genetics and Biology of Intraductal Papillary Mucinous Neoplasm. *Clin. Transl. Gastroenterol.* 2017, 8, e86.
7. Kopp, J.L.; Von Figura, G.; Mayes, E.; Liu, F.-F.; Dubois, C.L.; Morris, J.P.; Pan, F.C.; Akiyama, H.; Wright, C.V.; Jensen, K.; et al. Identification of Sox9-Dependent Acinar-to-Ductal Reprogramming as the Principal Mechanism for Initiation of Pancreatic Ductal Adenocarcinoma. *Cancer Cell* 2012, 22, 737–750.
8. Wagner, M.; Lührs, H.; Klöppel, G.; Adler, G.; Schmid, R.M. Malignant transformation of duct-like cells originating from acini in transforming growth factor α transgenic mice. *Gastroenterology* 1998, 115, 1254–1262.
9. Zhang, L.; Sanagapalli, S.; Stoita, A. Challenges in diagnosis of pancreatic cancer. *World J. Gastroenterol.* 2018, 24, 2047–2060.
10. Martin, G.R. Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc. Natl. Acad. Sci. USA* 1981, 78, 7634–7638.
11. Evans, M.J.; Kaufman, M.H. Establishment in culture of pluripotential cells from mouse embryos. *Nat. Cell Biol.* 1981, 292, 154–156.
12. Takahashi, K.; Yamanaka, S. Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* 2006, 126, 663–676.
13. González, F.; Boue, S.; Belmonte, J.C.I. Methods for making induced pluripotent stem cells: Reprogramming à la carte. *Nat. Rev. Genet.* 2011, 12, 231–242.
14. Malik, N.; Rao, M.S. A Review of the Methods for Human iPSC Derivation. *Methods Mol. Biol.* 2013, 997, 23–33.
15. Takahashi, K.; Tanabe, K.; Ohnuki, M.; Narita, M.; Ichisaka, T.; Tomoda, K.; Yamanaka, S. Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined Factors. *Cell* 2007, 131, 861–872.
16. Yu, J.; Vodyanik, M.A.; Smuga-Otto, K.; Antosiewicz-Bourget, J.; Frane, J.L.; Tian, S.; Nie, J.; Jonsdottir, G.A.; Ruotti, V.; Stewart, R.; et al. Induced Pluripotent Stem Cell Lines Derived from Human Somatic Cells. *science* 2007, 318, 1917–1920.
17. Shao, L.; Feng, W.; Sun, Y.; Bai, H.; Liu, J.; Currie, C.; Kim, J.; Gama, R.; Wang, Z.; Qian, Z.; et al. Generation of iPS cells using defined factors linked via the self-cleaving 2A sequences in a single open reading frame. *Cell Res.* 2009, 19, 296–306.

18. Fusaki, N.; Ban, H.; Nishiyama, A.; Saeki, K.; Hasegawa, M. Efficient induction of transgene-free human pluripotent stem cells using a vector based on Sendai virus, an RNA virus that does not integrate into the host genome. *Proc. Jpn. Acad. Ser. B* 2009, 85, 348–362.
19. Nishimura, K.; Sano, M.; Ohtaka, M.; Furuta, B.; Umemura, Y.; Nakajima, Y.; Ikehara, Y.; Kobayashi, T.; Segawa, H.; Takayasu, S.; et al. Development of Defective and Persistent Sendai Virus Vector. *J. Biol. Chem.* 2011, 286, 4760–4771.
20. Woltjen, K.; Michael, I.; Mohseni, P.; Desai, R.; Mileikovsky, M.; Hämäläinen, R.; Cowling, R.; Wang, W.; Liu, P.; Gertsenstein, M.; et al. piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells. *Nat. Cell Biol.* 2009, 458, 766–770.
21. Anokye-Danso, F.; Trivedi, C.; Jühr, D.; Gupta, M.; Cui, Z.; Tian, Y.; Zhang, Y.; Yang, W.; Gruber, P.J.; Epstein, J.A.; et al. Highly Efficient miRNA-Mediated Reprogramming of Mouse and Human Somatic Cells to Pluripotency. *Cell Stem Cell* 2011, 8, 376–388.
22. Warren, L.; Manos, P.D.; Ahfeldt, T.; Loh, Y.-H.; Li, H.; Lau, F.; Ebina, W.; Mandal, P.; Smith, Z.D.; Meissner, A.; et al. Highly Efficient Reprogramming to Pluripotency and Directed Differentiation of Human Cells with Synthetic Modified mRNA. *Cell Stem Cell* 2010, 7, 618–630.
23. Kim, D.; Kim, C.-H.; Moon, J.-I.; Chung, Y.-G.; Chang, M.-Y.; Han, B.-S.; Ko, S.; Yang, E.; Cha, K.Y.; Lanza, R.; et al. Generation of Human Induced Pluripotent Stem Cells by Direct Delivery of Reprogramming Proteins. *Cell Stem Cell* 2009, 4, 472–476.
24. Zhou, H.; Wu, S.; Joo, J.Y.; Zhu, S.; Han, D.W.; Lin, T.; Trauger, S.; Bien, G.; Yao, S.; Zhu, Y.; et al. Generation of Induced Pluripotent Stem Cells Using Recombinant Proteins. *Cell Stem Cell* 2009, 4, 381–384.
25. Okita, K.; Matsumura, Y.; Sato, Y.; Okada, A.; Morizane, A.; Okamoto, S.; Hong, H.; Nakagawa, M.; Tanabe, K.; Tezuka, K.-I.; et al. A more efficient method to generate integration-free human iPS cells. *Nat. Methods* 2011, 8, 409–412.
26. Okita, K.; Nakagawa, M.; Hyenjong, H.; Ichisaka, T.; Yamanaka, S. Generation of Mouse Induced Pluripotent Stem Cells Without Viral Vectors. *science* 2008, 322, 949–953.
27. Bock, C.; Kiskinis, E.; Verstappen, G.; Gu, H.; Boulting, G.; Smith, Z.D.; Ziller, M.; Croft, G.; Amoroso, M.W.; Oakley, D.; et al. Reference Maps of Human ES and iPS Cell Variation Enable High-Throughput Characterization of Pluripotent Cell Lines. *Cell* 2011, 144, 439–452.
28. Mallon, B.S.; Hamilton, R.S.; Kozhich, O.A.; Johnson, K.R.; Fann, Y.C.; Rao, M.S.; Robey, P. Comparison of the molecular profiles of human embryonic and induced pluripotent stem cells of isogenic origin. *Stem Cell Res.* 2014, 12, 376–386.
29. Mallon, B.S.; Chenoweth, J.G.; Johnson, K.R.; Hamilton, R.S.; Tesar, P.J.; Yavatkar, A.S.; Tyson, L.J.; Park, K.; Chen, K.; Fann, Y.C.; et al. StemCellDB: The Human Pluripotent Stem Cell Database at the National Institutes of Health. *Stem Cell Res.* 2013, 10, 57–66.

30. Choi, J.; Lee, S.; Mallard, W.; Clement, K.; Tagliazucchi, G.M.; Lim, H.; Choi, I.Y.; Ferrari, F.; Tsankov, A.M.; Pop, R.; et al. A comparison of genetically matched cell lines reveals the equivalence of human iPSCs and ESCs. *Nat. Biotechnol.* 2015, 33, 1173–1181.
31. Breunig, M.; Merkle, J.; Wagner, M.; Melzer, M.K.; Barth, T.F.; Engleitner, T.; Krumm, J.; Wiedenmann, S.; Cohrs, C.M.; Perkhofer, L.; et al. Modeling plasticity and dysplasia of pancreatic ductal organoids derived from human pluripotent stem cells. *Cell Stem Cell* 2021, 28, 1105–1124.
32. Calle, A.S.; Nair, N.; Oo, A.K.K.; Prieto-Vila, M.; Koga, M.; Khayrani, A.C.; Seno, M. A New PDAC Mouse Model Originated from iPSCs-Converted Pancreatic Cancer Stem Cells (CSCcm). *Am. J. Cancer Res.* 2016, 6, 2799–2815.
33. Huang, L.; Desai, R.; Conrad, D.N.; Leite, N.C.; Akshinthala, D.; Lim, C.M.; Gonzalez, R.; Muthuswamy, L.B.; Gartner, Z.; Muthuswamy, S.K. Commitment and oncogene-induced plasticity of human stem cell-derived pancreatic acinar and ductal organoids. *Cell Stem Cell* 2021, 28, 1090–1104.
34. Kim, J.; Hoffman, J.P.; Alpaugh, R.K.; Rhim, A.D.; Reichert, M.; Stanger, B.Z.; Furth, E.E.; Sepulveda, A.R.; Yuan, C.-X.; Won, K.J.; et al. An iPSC Line from Human Pancreatic Ductal Adenocarcinoma Undergoes Early to Invasive Stages of Pancreatic Cancer Progression. *Cell Rep.* 2013, 3, 2088–2099.
35. Hassan, G.; Afify, S.M.; Nair, N.; Kumon, K.; Osman, A.; Du, J.; Mansour, H.; Abu Quora, H.; Nawara, H.M.; Satoh, A.; et al. Hematopoietic Cells Derived from Cancer Stem Cells Generated from Mouse Induced Pluripotent Stem Cells. *Cancers* 2019, 12, 82.
36. Matsuda, S.; Yan, T.; Mizutani, A.; Sota, T.; Hiramoto, Y.; Prieto-Vila, M.; Chen, L.; Satoh, A.; Kudoh, T.; Kasai, T.; et al. Cancer stem cells maintain a hierarchy of differentiation by creating their niche. *Int. J. Cancer* 2014, 135, 27–36.
37. Nair, N.; Calle, A.S.; Zahra, M.H.; Prieto-Vila, M.; Oo, A.; Hurley, L.; Vaidyanath, A.; Seno, A.; Masuda, J.; Iwasaki, Y.; et al. A cancer stem cell model as the point of origin of cancer-associated fibroblasts in tumor microenvironment. *Sci. Rep.* 2017, 7, 1–13.
38. Cichocki, F.; Bjordahl, R.; Gaidarova, S.; Mahmood, S.; Abujarour, R.; Wang, H.; Tuininga, K.; Felices, M.; Davis, Z.B.; Bendzick, L.; et al. iPSC-derived NK cells maintain high cytotoxicity and enhance in vivo tumor control in concert with T cells and anti-PD-1 therapy. *Sci. Transl. Med.* 2020, 12, 1–15.
39. Deuse, T.; Hu, X.; Gravina, A.; Wang, D.; Tediashvili, G.; De, C.; Thayer, W.O.; Wahl, A.; Garcia, J.V.; Reichenspurner, H.; et al. Hypoimmunogenic derivatives of induced pluripotent stem cells evade immune rejection in fully immunocompetent allogeneic recipients. *Nat. Biotechnol.* 2019, 37, 252–258.

40. Klichinsky, M.; Ruella, M.; Shestova, O.; Lu, X.M.; Best, A.; Zeeman, M.; Schmierer, M.; Gabrusiewicz, K.; Anderson, N.R.; Petty, N.; et al. Human chimeric antigen receptor macrophages for cancer immunotherapy. *Nat. Biotechnol.* 2020, 38, 947–953.
41. Kooreman, N.G.; Kim, Y.; de Almeida, P.E.; Termglinchan, V.; Diecke, S.; Shao, N.-Y.; Wei, T.-T.; Yi, H.; Dey, D.; Nelakanti, R.; et al. Autologous iPSC-Based Vaccines Elicit Anti-tumor Responses In Vivo. *Cell Stem Cell* 2018, 22, 501–513.
42. Li, Y.; Hermanson, D.L.; Moriarity, B.S.; Kaufman, D.S. Human iPSC-Derived Natural Killer Cells Engineered with Chimeric Antigen Receptors Enhance Anti-tumor Activity. *Cell Stem Cell* 2018, 23, 181–192.e5.
43. Lu, S.; Zhang, Z.; Du, P.; Chard, L.S.; Yan, W.; El Khouri, M.; Wang, Z.; Zhang, Z.; Chu, Y.; Gao, D.; et al. A Virus-Infected, Reprogrammed Somatic Cell-Derived Tumor Cell (VIREST) Vaccination Regime Can Prevent Initiation and Progression of Pancreatic Cancer. *Clin. Cancer Res.* 2020, 26, 465–476.
44. Ouyang, X.; Liu, Y.; Zhou, Y.; Guo, J.; Wei, T.-T.; Liu, C.; Lee, B.; Chen, B.; Zhang, A.; Casey, K.M.; et al. Antitumor effects of iPSC-based cancer vaccine in pancreatic cancer. *Stem Cell Rep.* 2021, 16, 1468–1477.
45. Themeli, M.; Kloss, C.C.; Ciriello, G.; Fedorov, V.D.; Perna, F.; Gonen, M.; Sadelain, M. Generation of tumor-targeted human T lymphocytes from induced pluripotent stem cells for cancer therapy. *Nat. Biotechnol.* 2013, 31, 928–933.
46. Zhang, L.; Tian, L.; Dai, X.; Yu, H.; Wang, J.; Lei, A.; Zhu, M.; Xu, J.; Zhao, W.; Zhu, Y.; et al. Pluripotent stem cell-derived CAR-macrophage cells with antigen-dependent anti-cancer cell functions. *J. Hematol. Oncol.* 2020, 13, 1–5.
47. Rubio-Viqueira, B.; Jimeno, A.; Cusatis, G.; Zhang, X.; Iacobuzio-Donahue, C.; Karikari, C.; Shi, C.; Danenberg, K.; Danenberg, P.V.; Kuramochi, H.; et al. An In vivo Platform for Translational Drug Development in Pancreatic Cancer. *Clin. Cancer Res.* 2006, 12, 4652–4661.
48. Kim, M.P.; Evans, D.B.; Wang, H.; Abbruzzese, J.L.; Fleming, J.B.; E Gallick, G. Generation of orthotopic and heterotopic human pancreatic cancer xenografts in immunodeficient mice. *Nat. Protoc.* 2009, 4, 1670–1680.
49. Hermann, P.C.; Huber, S.L.; Herrler, T.; Aicher, A.; Ellwart, J.W.; Guba, M.; Bruns, C.J.; Heeschen, C. Distinct Populations of Cancer Stem Cells Determine Tumor Growth and Metastatic Activity in Human Pancreatic Cancer. *Cell Stem Cell* 2007, 1, 313–323.
50. Ishizawa, K.; Rasheed, Z.A.; Karisch, R.; Wang, Q.; Kowalski, J.; Susky, E.; Pereira, K.; Karamboulas, C.; Moghal, N.; Rajeshkumar, N.; et al. Tumor-Initiating Cells Are Rare in Many Human Tumors. *Cell Stem Cell* 2010, 7, 279–282.

51. Li, C.; Heidt, D.G.; Dalerba, P.; Burant, C.F.; Zhang, L.; Adsay, V.; Wicha, M.; Clarke, M.F.; Simeone, D.M. Identification of Pancreatic Cancer Stem Cells. *Cancer Res.* 2007, 67, 1030–1037.
52. Chen, L.; Kasai, T.; Li, Y.; Sugii, Y.; Jin, G.; Okada, M.; Vaidyanath, A.; Mizutani, A.; Satoh, A.; Kudoh, T.; et al. A Model of Cancer Stem Cells Derived from Mouse Induced Pluripotent Stem Cells. *PLoS ONE* 2012, 7, e33544.
53. Bailey, P.; Initiative, A.P.C.G.; Chang, D.K.; Nones, K.; Johns, A.L.; Patch, A.-M.; Gingras, M.-C.; Miller, D.K.; Christ, A.N.; Bruxner, T.J.C.; et al. Genomic analyses identify molecular subtypes of pancreatic cancer. *Nat. Cell Biol.* 2016, 531, 47–52.
54. Biankin, A.V.; Initiative, A.P.C.G.; Waddell, N.; Kassahn, K.S.; Gingras, M.-C.; Muthuswamy, L.B.; Johns, A.L.; Miller, D.K.; Wilson, P.J.; Patch, A.-M.; et al. Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. *Nat. Cell Biol.* 2012, 491, 399–405.
55. Witkiewicz, A.K.; McMillan, E.A.; Balaji, U.; Baek, G.; Lin, W.-C.; Mansour, J.C.; Mollaei, M.; Wagner, K.-U.; Koduru, P.; Yopp, A.C.; et al. Whole-exome sequencing of pancreatic cancer defines genetic diversity and therapeutic targets. *Nat. Commun.* 2015, 6, e6744.
56. Tu, Q.; Hao, J.; Zhou, X.; Yan, L.; Dai, H.; Sun, B.; Yang, D.; An, S.; Lv, L.; Jiao, B.; et al. CDKN2B deletion is essential for pancreatic cancer development instead of unmeaningful co-deletion due to juxtaposition to CDKN2A. *Oncogene* 2017, 37, 128–138.
57. Waddell, N.; Initiative, A.P.C.G.; Pajic, M.; Patch, A.-M.; Chang, D.K.; Kassahn, K.S.; Bailey, P.; Johns, A.L.; Miller, D.; Nones, K.; et al. Whole genomes redefine the mutational landscape of pancreatic cancer. *Nat. Cell Biol.* 2015, 518, 495–501.
58. Huang, L.; Holtzinger, A.; Jagan, I.; BeGora, M.; Lohse, I.; Ngai, N.; Nostro, M.C.; Wang, R.; Muthuswamy, L.B.; Crawford, H.C.; et al. Ductal pancreatic cancer modeling and drug screening using human pluripotent stem cell- and patient-derived tumor organoids. *Nat. Med.* 2015, 21, 1364–1371.
59. Hohwieler, M.; Illing, A.; Hermann, P.C.; Mayer, T.; Stockmann, M.; Perkhofer, L.; Eiseler, T.; Antony, J.S.; Müller, M.; Renz, S.; et al. Human pluripotent stem cell-derived acinar/ductal organoids generate human pancreas upon orthotopic transplantation and allow disease modelling. *Gut* 2016, 66, 473–486.
60. Ferreira, R.; Sancho, R.; Messal, H.A.; Nye, E.; Spencer-Dene, B.; Stone, R.K.; Stamp, G.; Rosewell, I.; Quaglia, A.; Behrens, A. Duct- and Acinar-Derived Pancreatic Ductal Adenocarcinomas Show Distinct Tumor Progression and Marker Expression. *Cell Rep.* 2017, 21, 966–978.
61. Zheng, X.; Carstens, J.; Kim, J.; Scheible, M.; Kaye, J.; Sugimoto, H.; Wu, C.-C.; LeBleu, V.S.; Kalluri, R. Epithelial-to-mesenchymal transition is dispensable for metastasis but induces

- chemoresistance in pancreatic cancer. *Nat. Cell Biol.* 2015, 527, 525–530.
62. Arumugam, T.; Ramachandran, V.; Fournier, K.F.; Wang, H.; Marquis, L.; Abbruzzese, J.L.; Gallick, G.E.; Logsdon, C.D.; McConkey, D.J.; Choi, W. Epithelial to Mesenchymal Transition Contributes to Drug Resistance in Pancreatic Cancer. *Cancer Res.* 2009, 69, 5820–5828.
63. Rhim, A.D.; Mirek, E.T.; Aiello, N.; Maitra, A.; Bailey, J.M.; McAllister, F.; Reichert, M.; Beatty, G.; Rustgi, A.K.; Vonderheide, R.H.; et al. EMT and Dissemination Precede Pancreatic Tumor Formation. *Cell* 2012, 148, 349–361.
64. Mitchem, J.; Brennan, D.J.; Knolhoff, B.L.; Belt, B.A.; Zhu, Y.; Sanford, D.E.; Belaygorod, L.; Carpenter, D.; Collins, L.; Piwnica-Worms, D.; et al. Targeting Tumor-Infiltrating Macrophages Decreases Tumor-Initiating Cells, Relieves Immunosuppression, and Improves Chemotherapeutic Responses. *Cancer Res.* 2013, 73, 1128–1141.
65. Lee, C.; Kozaki, T.; Ginhoux, F. Studying tissue macrophages in vitro: Are iPSC-derived cells the answer? *Nat. Rev. Immunol.* 2018, 18, 716–725.
66. Gutbier, S.; Wanke, F.; Dahm, N.; Rümmlin, A.; Zimmermann, S.; Christensen, K.; Köchl, F.; Rautanen, A.; Hatje, K.; Geering, B.; et al. Large-Scale Production of Human iPSC-Derived Macrophages for Drug Screening. *Int. J. Mol. Sci.* 2020, 21, 4808.
67. Von Ahrens, D.; Bhagat, T.D.; Nagrath, D.; Maitra, A.; Verma, A. The role of stromal cancer-associated fibroblasts in pancreatic cancer. *J. Hematol. Oncol.* 2017, 10, 1–8.
68. Ligorio, M.; Sil, S.; Malagon-Lopez, J.; Nieman, L.; Misale, S.; Di Pilato, M.; Ebright, R.Y.; Karabacak, N.M.; Kulkarni, A.S.; Liu, A.; et al. Stromal Microenvironment Shapes the Intratumoral Architecture of Pancreatic Cancer. *Cell* 2019, 178, 160–175.
69. Richards, K.E.; Zeleniak, A.E.; Fishel, M.; Wu, J.; Littlepage, L.E.; Hill, R. Cancer-associated fibroblast exosomes regulate survival and proliferation of pancreatic cancer cells. *Oncogene* 2017, 36, 1770–1778.
70. Sunami, Y.; Böker, V.; Kleeff, J. Targeting and Reprogramming Cancer-Associated Fibroblasts and the Tumor Microenvironment in Pancreatic Cancer. *Cancers* 2021, 13, 697.
71. Hosein, A.N.; Brekken, R.A.; Maitra, A. Pancreatic cancer stroma: An update on therapeutic targeting strategies. *Nat. Rev. Gastroenterol. Hepatol.* 2020, 17, 487–505.
72. Prieto-Vila, M.; Yan, T.; Calle, A.S.; Nair, N.; Hurley, L.; Kasai, T.; Seno, M. iPSC-Derived Cancer Stem Cells Provide a Model of Tumor Vasculature. *Am. J. Cancer Res.* 2016, 6, 1906–1921.
73. Yarchoan, M.; Hopkins, A.; Jaffee, E.M. Tumor Mutational Burden and Response Rate to PD-1 Inhibition. *N. Engl. J. Med.* 2017, 377, 2500–2501.
74. Brewer, B.G.; Mitchell, R.A.; Harandi, A.; Eaton, J.W. Embryonic vaccines against cancer: An early history. *Exp. Mol. Pathol.* 2009, 86, 192–197.

75. Ghosh, Z.; Huang, M.; Hu, S.; Wilson, K.D.; Dey, D.; Wu, J.C. Dissecting the Oncogenic and Tumorigenic Potential of Differentiated Human Induced Pluripotent Stem Cells and Human Embryonic Stem Cells. *Cancer Res.* 2011, 71, 5030–5039.
76. Ben-Porath, I.; Thomson, M.W.; Carey, V.J.; Ge, R.; Bell, G.W.; Regev, A.; Weinberg, R. An embryonic stem cell–like gene expression signature in poorly differentiated aggressive human tumors. *Nat. Genet.* 2008, 40, 499–507.
77. Zhao, T.; Zhang, Z.-N.; Rong, Z.; Xu, Y. Immunogenicity of induced pluripotent stem cells. *Nat. Cell Biol.* 2011, 13, 212–215.
78. De Almeida, P.E.; Meyer, E.H.; Kooreman, N.G.; Diecke, S.; Dey, D.; Sanchez-Freire, V.; Hu, S.; Ebert, A.D.; Odegaard, J.; Mordwinkin, N.M.; et al. Transplanted terminally differentiated induced pluripotent stem cells are accepted by immune mechanisms similar to self-tolerance. *Nat. Commun.* 2014, 5, 3903.
79. Tabatabai, R.; Linhares, Y.; Bolos, D.; Mita, M.; Mita, A. Targeting the Wnt Pathway in Cancer: A Review of Novel Therapeutics. *Target. Oncol.* 2017, 12, 623–641.
80. Argentiero, A.; De Summa, S.; Di Fonte, R.; Iacobazzi, R.M.; Porcelli, L.; Da Vià, M.; Brunetti, O.; Azzariti, A.; Silvestris, N.; Solimando, A.G. Gene Expression Comparison between the Lymph Node-Positive and -Negative Reveals a Peculiar Immune Microenvironment Signature and a Theranostic Role for WNT Targeting in Pancreatic Ductal Adenocarcinoma: A Pilot Study. *Cancers* 2019, 11, 942.
81. de Jesus, B.B.; Neves, B.M.; Ferreira, M.; Nóbrega-Pereira, S. Strategies for Cancer Immunotherapy Using Induced Pluripotent Stem Cells-Based Vaccines. *Cancers* 2020, 12, 3581.
82. Gąbka-Buszek, A.; Kwiatkowska-Borowczyk, E.; Jankowski, J.; Kozłowska, A.K.; Mackiewicz, A. Novel Genetic Melanoma Vaccines Based on Induced Pluripotent Stem Cells or Melanosphere-Derived Stem-Like Cells Display High Efficacy in a murine Tumor Rejection Model. *Vaccines* 2020, 8, 147.
83. Wang, J.; Shao, L.; Wu, L.; Ma, W.; Zheng, Y.; Hu, C.; Li, F. Expression levels of a gene signature in hiPSC associated with lung adenocarcinoma stem cells and its capability in eliciting specific antitumor immune-response in a humanized mice model. *Thorac. Cancer* 2020, 11, 1603–1612.
84. Li, Y.; Zeng, H.; Xu, R.-H.; Liu, B.; Li, Z. Vaccination with Human Pluripotent Stem Cells Generates a Broad Spectrum of Immunological and Clinical Response against Colon Cancer. *Stem Cells* 2009, 27, 3103–3111.
85. Lee, A.S.; Tang, C.; Rao, M.S.; Weissman, I.L.; Wu, J.C. Tumorigenicity as a clinical hurdle for pluripotent stem cell therapies. *Nat. Med.* 2013, 19, 998–1004.

86. Kooreman, N.G.; Wu, J.C. Tumorigenicity of pluripotent stem cells: Biological insights from molecular imaging. *J. R. Soc. Interface* 2010, 7, S753–S763.
87. Koyanagi-Aoi, M.; Ohnuki, M.; Takahashi, K.; Okita, K.; Noma, H.; Sawamura, Y.; Teramoto, I.; Narita, M.; Sato, Y.; Ichisaka, T.; et al. Differentiation-defective phenotypes revealed by large-scale analyses of human pluripotent stem cells. *Proc. Natl. Acad. Sci. USA* 2013, 110, 20569–20574.
88. Kawamura, A.; Miyagawa, S.; Fukushima, S.; Kawamura, T.; Kashiya, N.; Ito, E.; Watabe, T.; Masuda, S.; Toda, K.; Hatazawa, J.; et al. Teratocarcinomas Arising from Allogeneic Induced Pluripotent Stem Cell-Derived Cardiac Tissue Constructs Provoked Host Immune Rejection in Mice. *Sci. Rep.* 2016, 6, 19464.
89. Hochedlinger, K.; Yamada, Y.; Beard, C.; Jaenisch, R. Ectopic Expression of Oct-4 Blocks Progenitor-Cell Differentiation and Causes Dysplasia in Epithelial Tissues. *Cell* 2005, 121, 465–477.
90. Zhou, X.; Huang, G.-R.; Hu, P.; Xi, Z.; Guang-Rong, H.; Pin, H. Over-expression of Oct4 in human esophageal squamous cell carcinoma. *Mol. Cells* 2011, 32, 39–45.
91. Riggi, N.; Suvà, M.-L.; De Vito, C.; Provero, P.; Stehle, J.-C.; Baumer, K.; Cironi, L.; Janiszewska, M.; Petricevic, T.; Suvà, D.; et al. EWS-FLI-1 modulates miRNA145 and SOX2 expression to initiate mesenchymal stem cell reprogramming toward Ewing sarcoma cancer stem cells. *Genes Dev.* 2010, 24, 916–932.
92. Zhao, F.-Q.; Misra, Y.; Li, D.-B.; Wadsworth, M.P.; Krag, D.; Weaver, D.; Tessitore, J.; Zhang, G.; Tian, Q.; Buss, K. Differential expression of Oct3/4 in human breast cancer and normal tissues. *Int. J. Oncol.* 2018, 52, 2069–2078.
93. Bae, K.-M.; Su, Z.; Frye, C.; McClellan, S.; Allan, R.W.; Andrejewski, J.T.; Kelley, V.; Jorgensen, M.; Steindler, D.A.; Vieweg, J.; et al. Expression of Pluripotent Stem Cell Reprogramming Factors by Prostate Tumor Initiating Cells. *J. Urol.* 2010, 183, 2045–2053.
94. Sarkar, A.; Hochedlinger, K. The Sox Family of Transcription Factors: Versatile Regulators of Stem and Progenitor Cell Fate. *Cell Stem Cell* 2013, 12, 15–30.
95. Dang, C.V. MYC on the Path to Cancer. *Cell* 2012, 149, 22–35.
96. Clemente-Periván, S.I.; Gómez-Gómez, Y.; Leyva-Vázquez, M.A.; Lagunas-Martínez, A.; Organista-Nava, J.; Illades-Aguilar, B. Role of Oct3/4 in Cervical Cancer Tumorigenesis. *Front. Oncol.* 2020, 10, 247.
97. Themeli, M.; Rivière, I.; Sadelain, M. New Cell Sources for T Cell Engineering and Adoptive Immunotherapy. *Cell Stem Cell* 2015, 16, 357–366.

Retrieved from <https://encyclopedia.pub/entry/history/show/39526>